

3rd International Conference Advanced Mechanics: Structure, Materials, Tribology

Study of Non-Stationary Processes in Inhomogeneous Viscoelastic Materials

AIPCP25-CF-AMSMT2025-00021 | Article

PDF auto-generated using **ReView**



Study of Non-Stationary Processes in Inhomogeneous Viscoelastic Materials

Sergey G. Pshenichnov^{a)}, Ekaterina A. Korovaytseva^{b)} and
Anastasiya M. Razorenova^{c)}

Institute of Mechanics, Lomonosov Moscow State University, Michurinsky prospect 1, Moscow, Russia

^{a)} Corresponding author: serp56@yandex.ru

^{b)} katrell@mail.ru

^{c)} anastasija.razorenova@rambler.ru

Abstract. Initial boundary value problems that describe transient wave processes in inhomogeneous linearly viscoelastic bodies are investigated. A piecewise homogeneous structure with continuity conditions at the contact of homogeneous components is chosen as the main type of heterogeneity. At the same time, for viscoelastic functionally graded materials, their continuous heterogeneity is approximated by a layered homogeneous medium. The hereditary properties of homogeneous viscoelastic components are characterized by linear integral Boltzmann-Volterra constitutive relations with relaxation kernels of various types. In the mathematical formulation of the initial boundary value problem for a piecewise homogeneous viscoelastic body, the perturbations propagation area is considered limited. To construct solutions, the integral Laplace transform in time and the operation of its reversal are used. The solution in the originals is written in a new form. Special relations are presented that establish a certain correspondence between kernels belonging to different classes of functions. Using the example of a non-stationary problem for a functionally graded spherical layer, the similarity of non-stationary processes is demonstrated when using various singular kernels connected by these special relations.

Key words: inhomogeneous viscoelastic materials, hereditary kernels, Laplace transform, non-stationary waves, viscoelastic dynamic problems.

INTRODUCTION

The development of modern technologies, as well as the emergence of new materials, lead to the necessity of studying the dynamics of heterogeneous media with hereditary properties. These can be piecewise homogeneous structures with viscoelastic components, as well as viscoelastic functionally graded materials (FGM) with continuous spatial heterogeneity. Some results in this area, obtained recently by various authors for piecewise homogeneous materials, are contained, for example, in works [1-5]. At the same time, it should be noted that in most well-known publications, attention is paid not to transient wave processes, but to harmonic vibrations and waves [6-8]. In the field of viscoelastic FGM dynamics, other authors' research is devoted to stationary processes (for example, [9, 10]). Only some characteristic publications on the topic are mentioned here. Among the various approaches to studying non-stationary waves in media of this type, the development of analytical and numerical analytical methods is very important, but today the successes achieved in this area are clearly insufficient. This is especially true for transient wave processes in piecewise homogeneous viscoelastic bodies with a large number of homogeneous components and in FGM.

This paper contains the results of research in the field of unsteady dynamics of viscoelastic inhomogeneous media within the framework of the Boltzmann-Volterra model, conducted by using numerical analytical methods based on previous theoretical developments [11 – 13]. The solutions of initial boundary value problems of viscoelasticity for piecewise homogeneous bodies are presented in a special form that is suitable for different types of relaxation kernels.

When using linear integral Boltzmann-Volterra constitutive relations, the question arises as to how belonging of viscoelastic kernels to one or another class of functions affects the transition process. This issue has already been raised earlier [11, 14, 15], and here the proposed approach has been further developed. Using the example of a non-stationary problem for a spherical FGM layer, the similarity of non-stationary processes is demonstrated when using various singular kernels in the Boltzmann-Volterra model, connected by special relations.

FORMULATION OF THE NON-STATIONARY PROBLEM. THE SOLUTION REPRESENTATION FORM

The layered homogeneous medium with continuity conditions at the contact of the layers is considered the main type of heterogeneity. The continuous heterogeneity of viscoelastic FGMs will be approximated by a structure consisting of a large number of viscoelastic homogeneous layers. This approach is widely used in modeling stationary processes in elastic, thermoelastic, and piezoelectroelastic FGMs [16 – 18]. In non-stationary problems, the validity of such an approximation is confirmed by calculations for various bodies (see, for example, [19]). Let us consider a nonstationary linear viscoelasticity problem for the N -component piecewise homogeneous body, including the equations of dynamics ($n = 1, 2, \dots, N$):

$$(\hat{\lambda}_n + \hat{\mu}_n) \text{grad div } \mathbf{u}^{(n)}(\mathbf{x}, t) + \hat{\mu}_n \Delta \mathbf{u}^{(n)}(\mathbf{x}, t) + \mathbf{f}^{(n)}(\mathbf{x}, t) = \rho_n \ddot{\mathbf{u}}^{(n)}(\mathbf{x}, t), \quad \mathbf{x} \in \Omega_n, \quad (1)$$

constitutive relations:

$$\tilde{\boldsymbol{\sigma}}^{(n)}(\mathbf{x}, t) = 2\hat{\mu}_n \tilde{\boldsymbol{\varepsilon}}^{(n)}(\mathbf{x}, t) + \hat{\lambda}_n \text{div } \mathbf{u}^{(n)}(\mathbf{x}, t) \tilde{\mathbf{I}}, \quad \tilde{\boldsymbol{\varepsilon}}^{(n)} = \text{def } \mathbf{u}^{(n)}, \quad \mathbf{x} \in \Omega_n, \quad (2)$$

boundary conditions (m is the number of the component that has common points with the boundary Σ):

$$\tilde{\boldsymbol{\alpha}}^{(m)}(\mathbf{x}) \tilde{\boldsymbol{\sigma}}^{(m)}(\mathbf{x}, t) \mathbf{n} + \tilde{\boldsymbol{\beta}}^{(m)}(\mathbf{x}) \mathbf{u}^{(m)}(\mathbf{x}, t) = \mathbf{p}^{(m)}(\mathbf{x}, t), \quad \mathbf{x} \in \Sigma, \quad t > 0, \quad (3)$$

continuity conditions at the contact of adjacent components with numbers p and q :

$$\mathbf{u}^{(p)}(\mathbf{x}, t) = \mathbf{u}^{(q)}(\mathbf{x}, t), \quad \tilde{\boldsymbol{\sigma}}^{(p)}(\mathbf{x}, t) \mathbf{n} = \tilde{\boldsymbol{\sigma}}^{(q)}(\mathbf{x}, t) \mathbf{n}, \quad \mathbf{x} \in \Sigma_{pq}, \quad (4)$$

and initial conditions:

$$\mathbf{u}^{(n)}(\mathbf{x}, 0) = \mathbf{b}_1^{(n)}(\mathbf{x}), \quad \dot{\mathbf{u}}(\mathbf{x}, 0) = \mathbf{b}_2^{(n)}(\mathbf{x}), \quad \mathbf{x} \in \Omega_n, \quad (5)$$

herewith the operators $\hat{\lambda}_n$ and $\hat{\mu}_n$ have the form

$$\hat{\lambda}_n = \frac{1}{3} [3K_0^{(n)} (1 - \hat{T}_v^{(n)}) - 2G_0^{(n)} (1 - \hat{T}_s^{(n)})], \quad \hat{\mu}_n = G_0^{(n)} (1 - \hat{T}_s^{(n)}), \quad \hat{T}_j^{(n)} \xi(t) = \int_0^t T_j^{(n)}(t - \tau) \xi(\tau) d\tau, \quad j = v, s \quad (6)$$

Here the indexes of the values correspond to the components numbers; $\tilde{\boldsymbol{\sigma}}^{(n)}$, $\tilde{\boldsymbol{\varepsilon}}^{(n)}$ are the stress and small strain tensors; $\mathbf{u}^{(n)}$ is the displacement vector; $\mathbf{n}$ is an exterior unit normal to the corresponding boundary; ρ_n is the density; Δ is Laplace operator; $\tilde{\mathbf{I}}$ is the unit tensor; $G_0^{(n)}$, $K_0^{(n)}$ are the instantaneous values of the shear and bulk moduli; $T_v^{(n)}(t)$, $T_s^{(n)}(t)$ are the volumetric and shear relaxation kernels; $\tilde{\boldsymbol{\alpha}}^{(m)}$, $\tilde{\boldsymbol{\beta}}^{(m)}$ are given second-rank tensors that determine the type of boundary conditions; $\mathbf{p}^{(m)}$, $\mathbf{f}^{(n)}$, $\mathbf{b}_1^{(n)}$, $\mathbf{b}_2^{(n)}$ are given vectors of boundary effects, volumetric forces, initial displacements and velocities; the dot above the variable denotes the derivative with respect to time t . The disturbances propagations domain Ω consists from N subdomains: $\Omega = \Omega_1 \cup \Omega_2 \cup \dots \cup \Omega_N$, which correspond to homogeneous isotropic components of the body, herewith Ω and its boundary Σ are limited. This does not necessarily mean that the body itself is limited. The dimension of the coordinate vector $\mathbf{x}$ depends on the specific problem and can vary from 1 to 3.

The paper [11] contains the results of general theoretical studies of problem (1)–(6) based on the application of the integral Laplace transform with respect to time and the subsequent construction of a solution in the originals. It was established the relationship between the properties of the set E of eigenvalues of the free oscillations problem for the body under consideration and the branching points of the solution of the non-stationary problem (1) - (6) in the space of Laplace images, as well as the relationship between the elements of the set E and the poles of the solution in the images. In [12], new forms of the solution to the linear viscoelasticity problem for a homogeneous body are presented. For the case of a piecewise homogeneous viscoelastic medium considered here, similar mathematical considerations make it possible to write down the solution in a new form, taking into account the set of homogeneous components of the material.

So, let us apply the integral Laplace transform in time to the problem (1) – (6). Denote the images of the functions $\mathbf{u}^{(n)}(\mathbf{x}, t)$, $\tilde{\boldsymbol{\sigma}}^{(n)}(\mathbf{x}, t)$, $\mathbf{p}^{(n)}$, $\mathbf{f}^{(n)}$, $T_v^{(n)}(t)$, $T_s^{(n)}(t)$ as $\mathbf{U}^{(n)}(\mathbf{x}, s)$, $\tilde{\mathbf{S}}^{(n)}(\mathbf{x}, s)$, $\mathbf{P}^{(n)}$, $\mathbf{F}^{(n)}$, $\Theta_v^{(n)}(s)$, $\Theta_s^{(n)}(s)$ ($s \in C$) respectively.

Assume that there are no volumetric forces: $\mathbf{f}^{(n)} \equiv \mathbf{0}$, and for surface loads the following relation is satisfied:

$$\mathbf{p}^{(n)}(\mathbf{x}, t) = \mathbf{q}^{(n)}(\mathbf{x})\psi(t), \quad \psi(t) = h(t),$$

where $h(t)$ is the Heaviside function, and there are no displacements of the body as a rigid whole. Let herewith one of the following two cases be realized for the hereditary properties of a piecewise homogeneous body.

Case 1. All non-identically zero hereditary kernels $T_v^{(n)}$, $T_s^{(n)}$ belong to the class of functions that are partial sums of the Prony series:

$$\sum_{m=1}^M a_m \exp(-b_m t), \quad \text{where} \quad \sum_{m=1}^M a_m / b_m < 1, \quad a_m > 0, \quad b_m > 0 \quad (m = 1, 2, \dots, M), \quad (7)$$

Case 2. $T_v^{(n)} \equiv 0$ for every n , and all non-identically zero $T_s^{(n)}$ are Rzhantsyn-Koltunov kernels (here the time is assumed to be dimensionless):

$$A e^{-Bt} t^{\delta-1}, \quad \text{where} \quad A\Gamma(\delta)/B^\delta < 1, \quad A > 0, \quad B > 0, \quad 0 < \delta < 1, \quad (8)$$

$\Gamma(\delta)$ is a gamma-function with an appropriate argument. The constants a_m, b_m and M , as well as the constants A, B, δ for each component of a piecewise homogeneous medium, are, generally speaking, unique.

Besides, let the asymptotic behavior of the images $\mathbf{U}^{(n)}$ with $s \rightarrow \infty$ be “favorable” [12]. Then, reasoning similarly to how it was done in the work [12], we can represent the originals $\mathbf{u}^{(n)}$ in a form containing a series of residues at the complex poles z_j of the images, as well as two integrals (one over a finite segment, the other is improper):

$$\begin{aligned} \mathbf{u}^{(n)}(\mathbf{x}, t) = & \text{Res}_{s=0}[\mathbf{U}^{(n)}(\mathbf{x}, s)e^{st}] + 2 \sum_{j=1}^{\infty} \text{Re}\{\text{Res}_{s=z_j}[\mathbf{U}^{(n)}(\mathbf{x}, s)e^{st}]\} + \frac{1}{\pi} e^{\alpha_0 t} \int_0^{\alpha_0} \text{Re}\{\mathbf{U}^{(n)}(\mathbf{x}, \alpha_0 + i\omega)e^{i\omega t}\} d\omega - \\ & - \frac{1}{\pi} \int_{-\infty}^{\alpha_0} \text{Im}\{\mathbf{U}^{(n)}(\mathbf{x}, \alpha + i\omega_0)e^{i\omega_0 t}\} e^{\alpha t} d\alpha, \quad n = 1, 2, \dots, N. \end{aligned} \quad (9)$$

Here $z_j = \alpha_j + i\omega_j$, $\alpha_j, \omega_j \in R$, $\omega_j > 0$, $j = 1, 2, 3, \dots$, herewith $\alpha_j < 0$. The values α_0, ω_0 are chosen arbitrarily at intervals $\alpha_0 \in (s^*; 0)$, $\omega_0 \in (0; \omega_1)$, where $\omega_1 = \text{Im}(z_1)$, z_1 is the complex pole with the smallest positive imaginary part, s^* is the largest among the roots of all equations:

$$1 - \Theta^{(n)} = 0, \quad 1 - \Theta_s^{(n)} = 0, \quad (10)$$

where $\Theta_s^{(n)}(s)$ are Laplace images of kernels $T_s^{(n)}(t)$, and $\Theta^{(n)}$ are images of functions:

$$T^{(n)}(t) = [(1 + \nu_0^{(n)})T_v^{(n)}(t) + 2(1 - 2\nu_0^{(n)})T_s^{(n)}(t)] / [3(1 - \nu_0^{(n)})].$$

Note that $\{\alpha_j \pm i\omega_j\} \subset E$ and if a piecewise homogeneous body has at least one viscoelastic component, then the set E can have finite limit points on the complex plane. These points do not belong to E and are the roots of at least one of the equations (10). In the above two cases concerning the hereditary properties of the components, the roots of equations (10) are real negative ones.

The expression for the stress tensor $\tilde{\boldsymbol{\sigma}}^{(n)}$ through its image $\tilde{\mathbf{S}}^{(n)}$ is similar to (9). For other loading functions $\psi(t)$ solutions are obtained based on formula (9) and the Duhamel integral. The field of applicability of formula (9) is probably much broader than the indicated cases 1 and 2, but this issue requires further study.

SPECIAL RELATIONS BETWEEN RELAXATION KERNELS

When studying wave processes in linearly viscoelastic media, an important question arises about the choice of a class of functions for relaxation kernels $T_v^{(n)}, T_s^{(n)}$, which are included in the constitutive Boltzmann-Volterra relations (6). We introduce dimensionless hereditary kernels and time:

$$\gamma_s^{(n)}(\tau) = t_0 T_s^{(n)}, \quad \gamma_v^{(n)}(\tau) = t_0 T_v^{(n)}, \quad \tau = t / t_0, \quad (11)$$

where t_0 is some characteristic process time. Next, we will consider the dimensionless kernels $\gamma_s^{(n)}, \gamma_v^{(n)}$ in the class of functions (7) or (8), considering all constants and the time dimensionless in accordance with relations (11) and formally replacing t with τ in (7), (8). Note that the kernels from class (8) have a weak singularity at zero, unlike regular kernels (7). Consider the question of how significantly the belonging of relaxation kernels to class (7) or class (8) affects the nature of unsteady wave processes in a viscoelastic medium. In this regard, special relations have been proposed [11, 14, 15] that establish a certain correspondence between hereditary kernels from different classes of functions. Let us formulate these relations here using dimensionless quantities.

Let given a kernel $\gamma_1(\tau)$ (volumetric or shear relaxation) and chosen a class of kernels Q , defined by a known function $\Phi(\beta_1, \beta_2, \dots, \beta_K, \tau)$, which depends on K the numerical parameters $(\beta_1, \beta_2, \dots, \beta_K) \in \Omega_\beta \subset \square^K$ and the time τ . We construct the kernel $\gamma_2(\tau) = \Phi(\beta_1^0, \beta_2^0, \dots, \beta_K^0, \tau) \in Q$ by determining the values $\beta_1^0, \beta_2^0, \dots, \beta_K^0$ from the condition of equality of the long-term moduli corresponding to the kernels γ_1 and γ_2 :

$$\int_0^\infty \Phi(\beta_1, \beta_2, \dots, \beta_K, \tau) d\tau = \int_0^\infty \gamma_1(\tau) d\tau, \quad (12)$$

and minimizing the integral (if it exists) by the parameters β_j :

$$\int_0^\infty [\Phi(\beta_1, \beta_2, \dots, \beta_K, \tau) - \gamma_1(\tau)]^2 d\tau \xrightarrow{\beta_1, \beta_2, \dots, \beta_K} \min, \quad (13)$$

It was hypothesized that the kernels $\gamma_1(\tau)$ and $\gamma_2(\tau)$, connected by relations (12), (13), affect transient processes in viscoelastic materials similarly, all other initial data being equal, and some possible restrictions on the domain Ω_β of variation of the parameters β_j .

Consider as a class Q the set of functions of the form (7) with one term of the sum ($M=1$), i.e., with only two parameters: $\beta_1 = a_1$, $\beta_2 = b_1$. It was proved [15] that if the initial kernel $\gamma_1(\tau)$ is a smooth function and $[\gamma_1(\tau)]^2$ is integrable on $(0; \infty)$, and the following conditions are satisfied:

$$\lim_{\tau \rightarrow +\infty} \tau \gamma_1(\tau) = 0, \quad \lim_{\tau \rightarrow 0+} \tau \gamma_1(\tau) = 0, \quad d\gamma_1(\tau)/d\tau < 0 \quad (\tau > 0), \quad (14)$$

then for $\gamma_1(\tau)$ there exists, moreover, a single kernel $\gamma_2(\tau) \in Q$:

$$\gamma_2(\tau) = \beta_1^0 \exp(-\beta_2^0 \tau), \quad (15)$$

satisfying conditions (12), (13). Herewith $\beta_1^0 = \beta_2^0 \Theta_1(0)$, and β_2^0 is the only positive root of the algebraic equation (with respect to the unknown β):

$$\Theta_1(0) - 4[\Theta_1(\beta) + \beta \Theta_1'(\beta)] = 0, \quad \beta > 0, \quad (16)$$

where $\Theta_1(s)$ is the Laplace transform of $\gamma_1(\tau)$; the prime denotes the derivative. Note that if the initial kernel γ_1 is chosen from the set of functions (7) with $M \geq 2$, or (8) with the additional requirement $0.5 < \delta < 1$, then its square is integrable on $(0; \infty)$ and it has all the properties (14).

The hypothesis about the similarity of the influence of the corresponding hereditary kernels was confirmed by calculations for a number of specific examples [11, 14, 15], where the initial kernel $\gamma_1(\tau)$ was chosen either in the form (7) for $M \geq 2$, or in the form (8) for $0.5 < \delta < 1$, and the kernel $\gamma_2(\tau)$ was searched for in the two-parameter form (15). If the singularity parameter for the initial kernel γ_1 was chosen not too close to 0.5, namely: $0.6 \leq \delta < 1$, then in all examples the similarity of non-stationary processes was observed for the kernels $\gamma_1(\tau)$ and $\gamma_2(\tau)$. This similarity was assessed visually by graphs of the dependence of stresses and displacements on the time at various fixed points of the bodies. The dynamics of some homogeneous bodies [14, 15], as well as the dynamics of an infinite two-layer cylinder [11], were investigated. In the present work, relying on previous results, we consider the influence of singular relaxation kernels of the form (8) on transient processes in viscoelastic FGM. Herewith the parameter δ is varied, and the singular kernels themselves are connected by certain relations.

So, let the singular relaxation kernel γ_1 have the form (8). We construct a two-parameter kernel γ_2 (15) for it using relations (12), (13). Taking into account the equation (16), as well as the type of the kernel γ_1 and its image $\Theta_1(s)$, we obtain the following relations for the quantities β_1^0, β_2^0 :

$$\beta_2^0 = B[y_0(\eta) - 1], \quad \beta_1^0 = \beta_2^0 A \Gamma(\delta) / B^\delta, \quad (17)$$

where $\eta = 1 - \delta$, $0 < \eta < 0.5$, herewith $y_0(\eta)$ is the root of an algebraic equation (with a parameter η) relatively unknown y :

$$y^{2-\kappa} - 4\eta y - 4(1-\eta) = 0.$$

For a given η ($0 < \eta < 0.5$), this root exists and is the only one, and $y_0(\eta) > 1$. The second relation (17) means the equality of the long-term moduli (12) corresponding to the kernels γ_1 and γ_2 , and the first is obtained by minimizing the integral (13) by two parameters: β_1 and β_2 .

Let us rewrite the relations (17) for the three parameters of the kernel γ_1 in the form

$$B = \beta_2^0 / [y_0(\eta) - 1], \quad A = \beta_1^0 B^\delta / [\beta_2^0 \Gamma(\delta)]. \quad (18)$$

Thus, a regular two-parameter kernel γ_2 of the form (15) corresponds to an infinite set of singular three-parameter kernels γ_1 of the form (8). Choosing, for example, a certain value of the singularity parameter δ , one can easily find the remaining two parameters A, B of the singular kernel (8), which is in accordance with the kernel of the form (15) according to the relations (12), (13). Next, using a specific example of a spherical FGM layer, we will demonstrate the effect of hereditary kernels of the form (8) with various singularity parameters on transient wave processes. In this case, such singular kernels are in accordance with the same regular two-parameter kernel (15) according to the relations (18).

CALCULATION RESULTS FOR A SPHERICAL FGM LAYER

In a spherical coordinate system R, θ, ϕ , let us consider the problem of the unsteady longitudinal waves propagation in a spherical layer (hollow sphere) consisting of a viscoelastic FGM. The instantaneous values of the shear modulus G_0 and Poisson's ratio ν_0 , as well as the density ρ of this material, generally speaking, continuously depend on the radial coordinate R . In addition, we will consider the kernel of the shear relaxation T_s to be a function of R and t , and the kernel of the volumetric relaxation T_v to be identically zero. Let's denote the inner and outer radii of the layer by R_0 and R_{sph} , accordingly. The initial state of the body is rest, but from the moment $t = 0$ a uniformly distributed normal load $P(t)$ begins to act on its outer surface $R = R_{\text{sph}}$, while the inner surface $R = R_0$ remains free (Fig. 1).




FIGURE 1. The central section of the FGM sphere and the loading scheme

We approximate the continuous heterogeneity of the FGM by concentric homogeneous viscoelastic spherical layers with kinematic and dynamic continuity conditions at the boundaries between them. The relations $R_{n-1} \leq R \leq R_n$, $n = 1, 2, 3, \dots, N$ correspond to the layers, herewith $R_N = R_{\text{sph}}$, $N \gg 1$. Let us introduce dimensionless values:

$$r = R / R_N, \quad r_0 = R_0 / R_N, \quad r_n = R_n / R_N \quad (n=1,2,3,\dots,N, \quad r_N=1), \quad \tau = t / t_0, \quad \psi(\tau) = P / F_0 \quad (19)$$

where $t_0 = R_N / c_N$, c_N is the longitudinal elastic wave velocity in the N -th layer; F_0 is the characteristic constant of the stress dimension. Dimensionless shear relaxation kernel $\gamma_s^{(n)}(\tau)$ has the form (8) taking into account the relations (11), in which it is accepted: $t_0 = R_N / c_N$. For the initial FGM layer also: $r_0 = R_0 / R_{sph} = R_0 / R_N$ and $\tau = t / t_0$, where $t_0 = R_{sph} / c(1)$, $c(r)$ is the longitudinal elastic wave velocity in the FGM. Note that the equality $c(1) = c_N$ is satisfied exactly or approximately, depending on the method of approximating the continuous heterogeneity by layers. The mathematical formulation of the spherically symmetric problem under consideration for a multilayer viscoelastic sphere is contained in [13]. This work also contains the solution to this problem in images after applying the integral Laplace transform in time and the solution in the originals based on formula (9).

Here we consider a spherical viscoelastic FGM layer with $r_0 = 0.8$ and the following initial data ($r_0 \leq r \leq 1$):

$$\frac{G_0(r)}{G_0(r_0)} = \frac{\rho(r)}{\rho(r_0)} = 1 + 4 \frac{r - r_0}{1 - r_0}, \quad \nu_0 \equiv 0.3; \quad \gamma_v \equiv 0, \quad \gamma_s(r, \tau) = a(r)e^{-\tau}, \quad a(r) = 0.3(1 - r) / (1 - r_0). \quad (20)$$

It follows from relations (20) that the velocity of longitudinal elastic waves c does not depend on r , which is convenient when analyzing transients. The kernel γ_s has a form similar to (15), but with the coefficient $\beta_1^0 = a(r)$ that is not a constant, but a function that decreases linearly with respect to r . The values G_0, ρ increase linearly over r , herewith $G_0(1) / G_0(r_0) = \rho(1) / \rho(r_0) = 5$. We choose the function characterizing the load in the form of a smoothed step: $\psi(\tau) = 1 - e^{-50\tau}$ ($\tau > 0$) and then the load P limit with $\tau \rightarrow \infty$ is equal to F_0 (19). During the research were found the values of relative stresses:

$$\kappa_1(r, \tau) = \sigma_R / F_0, \quad \kappa_2(r, \tau) = \sigma_\theta / F_0,$$

where σ_R, σ_θ are radial and circumferential stresses ($\sigma_\theta = \sigma_\phi$). They were approximated by the corresponding values for a layered homogeneous structure:

$$\kappa_1^{(n)}(r, \tau) = \sigma_R^{(n)} / F_0, \quad \kappa_2^{(n)}(r, \tau) = \sigma_\theta^{(n)} / F_0, \quad n = 1, 2, 3, \dots, N,$$

herewith, the parameters of the FGM were approximated by 80 layers of equal thickness (this turned out to be quite sufficient). All calculations were carried out based on the solution of the problem for the corresponding multilayer hollow sphere [13] using formula (9).

For the given shear relaxation kernel (20), we introduce the notation $\gamma_0 = \gamma_s(r, \tau)$ and, for this kernel, according to relations (18), we find the parameters A, B of the Rzhantsyn-Koltunov singular kernel (8) by specifying different values of the singularity parameter δ . Here, B is a constant, and A is a linearly decreasing function of the variable r . We denote the singular shear relaxation kernels constructed in this way for three different values of the parameter δ by $\gamma_1, \gamma_2, \gamma_3$ and present the expressions obtained for them:

$$\begin{aligned} \delta = 0.9 \quad (\eta = 0.1), \quad \gamma_1(r, \tau) &= A(r)e^{-0.832\tau}\tau^{-0.1}, \quad A(r) = 0.238(1 - r) / (1 - r_0), \\ \delta = 0.7 \quad (\eta = 0.3), \quad \gamma_2(r, \tau) &= A(r)e^{-0.508\tau}\tau^{-0.3}, \quad A(r) = 0.144(1 - r) / (1 - r_0), \\ \delta = 0.55 \quad (\eta = 0.45), \quad \gamma_3(r, \tau) &= A(r)e^{-0.285\tau}\tau^{-0.45}, \quad A(r) = 0.093(1 - r) / (1 - r_0). \end{aligned}$$

Figures 2–4 show the time dependences of the relative circumferential stress κ_2 at different fixed points of the FGM layer for kernels $\gamma_0, \gamma_1, \gamma_2, \gamma_3$ (curves 1–4, respectively) with all other initial data (20) being identical. The figures also present the results for a spherical elastic FGM layer (curves 5) and for a homogeneous elastic spherical layer (curves 6) of the same dimensions. Figure 5 shows similar results for the relative radial stress κ_1 . Everywhere negative stresses are compressive.




FIGURE 2. Evolution of κ_2 in time at the point $r = r_0 = 0.8$ (1 – with the regular kernel γ_0 ; 2 – 4 – with the singular kernels $\gamma_1, \gamma_2, \gamma_3$; 5 – for the elastic FGM; 6 – for the homogeneous elastic spherical layer)




FIGURE 3. Evolution of κ_2 in time at the point $r = 1$




FIGURE 4. Evolution of κ_2 in time at the point $r = 0.9$




FIGURE 5. Evolution of κ_1 in time at the point $r = 0.9$

In the considered time range for viscoelastic FGMs, the similarity of nonstationary processes to each other is generally observed for all four shear relaxation kernels $\gamma_0, \gamma_1, \gamma_2, \gamma_3$. Curves 1 and 2 practically merge with each other. It should be noted that, as time progresses, the transient process for kernel γ_3 differs somewhat more from the processes for the other hereditary kernels. The figures also demonstrate the influence of viscosity and material inhomogeneity on the transient wave process.

CONCLUSION

Based on the integral Laplace transform in time and its reversal, new form of representation of solutions to non-stationary problems for inhomogeneous viscoelastic bodies is obtained within the framework of the Boltzmann-Volterra model for various types of hereditary kernels. This makes it possible in many cases to significantly simplify the dynamic calculations of various structures.

The influence of singular Rzhantsyn-Koltunov relaxation kernels on transient processes in spherical viscoelastic FGM layer was investigated. It was found that hereditary kernels, related by the proposed special relations, in general have a similar effect on transient processes in viscoelastic materials.

ACKNOWLEDGMENTS

The research was supported by the Russian Science Foundation, grant No 24-29-00164, <https://rscf.ru/en/project/24-29-00164/>.

REFERENCES

1. V.G. Bazhenov, A.A. Belov, L.A. Igumnov, *Boundary element modeling of the dynamics of piecewise homogeneous media and structures*, (NNGU, N. Novgorod, 2009), 180 p. [In Russian].
2. A.N. Petrov, M.D. Ermolaev, Boundary elements analysis of the dynamics of composite viscoelastic bodies. *Vestnik of Lobachevsky State University of Nizhny Novgorod. Solid Mechanics* **4**(4), 1694–1696 (2011), [In Russian].
3. Sh. Hosseini Hashemi and H. Bakhshi Khaniki, Dynamic behavior of multi-layered viscoelastic nanobeam system embedded in a viscoelastic medium with a moving nanoparticle. *J. Mech.* **33**(5), 559–575 (2017). <https://doi.org/10.1017/jmech.2016.91>
4. H.S. Lee, Viscowave – a new solution for viscoelastic wave propagation of layered structures subjected to an impact load. *Int. J. Pavement Eng.* **15**(6), 542 – 557 (2014). <https://doi.org/10.1080/10298436.2013.782401>
5. K. Khudoynazarov, Longitudinal-radial vibrations of a viscoelastic cylindrical three-layer structure. *Facta universitatis. Series: Mechanical Engineering. Special Issue.* **22**(3), 473 – 484 (2024). <https://doi.org/10.22190/FUME231219010K>

6. V. Alizadeh and A.V. Amirkhizi, Overall dynamic properties of locally resonant viscoelastic layered media based on consistent field integration for oblique anti-plane shear waves. *Mech. Mater.* **160**. Art. 103981 (2021). <https://doi.org/10.1016/j.mechmat.2021.103981>
7. R.D. Borchardt, *Viscoelastic Waves and Rays in Layered Media*, (Cambridge University Press, Cambridge, 2020), 450 p.
8. A.S. Shamaev, V.V. Shumilova, Asymptotics of the spectra of one-dimensional natural vibrations in media consisting of solid and fluid layers. *Dokl. Phys.* **65**(4), 153 – 156 (2020). <https://doi.org/10.1134/S1028335820040084>.
9. A.O. Vatulyan, V.O. Yurov, Waves in a viscoelastic cylindrical waveguide with a defect. *Izvestiya of Saratov University. Mathematics. Mechanics. Informatics* **21**(3), 352–367 (2021) [In Russian], <https://doi.org/10.18500/1816-9791-2021-21-3-352-367>
10. X. Zhang, Z. Li, X. Wang, J.G. Yu, The fractional Kelvin-Voigt model for circumferential guided waves in a viscoelastic FGM hollow cylinder. *Applied Mathematical Modelling* **89**. Part 1, 299-313 (2021). <https://doi.org/10.1016/j.apm.2020.06.077>
11. S.G. Pshenichnov, *Mechanics of Solids* **51**(1), 65–74 (2016). <https://doi.org/10.3103/S0025654416010076>
12. S. Pshenichnov, E. Korovaytseva, A. Razorenova, Study of transient wave processes in viscoelastic materials. *AIP Conf. Proc.* **3177**. Art 050007 (2025). <https://doi.org/10.1063/5.0294734>
13. S.G. Pshenichnov, Problems of Strength and Plasticity **87**(1), 103 - 112 (2025), [in Russian]. <https://doi.org/10.32326/1814-9146-2025-87-1-103-112>
14. E.A. Korovaytseva, S.G. Pshenichnov, T. Zhelyazov, M.D. Datcheva, On the problem of non-stationary waves propagation in a linear-viscoelastic layer. *Comptes Rendus de l'Academie Bulgare des Sciences* **74**(5), 748 – 755 (2021). <https://doi.org/10.7546/CRABS.2021.05.13>
15. S.G. Pshenichnov, M.Yu. Starovskaya. The equivalence conditions for relaxation kernels in nonstationary problems in viscous elasticity. *Izvestiya Of the Tula State University. Ser. Mathematics. Mechanics. Informatics* **12**(2), 177 – 189 (2006), [in Russian].
16. P. Kielczynski, M. Szalewski, A. Balcerzak, K. Wieja, Propagation of ultrasonic Love waves in nonhomogeneous elastic functionally graded materials. *Ultrasonics* **65**, 220-227 (2016). <https://doi.org/10.1016/j.ultras.2015.10.001>
17. N.V. Larin, L.A. Tolokonnikov, Sound scattering by a thermoelastic ball with a continuously inhomogeneous coating in a heat-conducting fluid. *Math. Model. Comput. Simul.* **11**(6), 1007–1018 (2019). <https://doi.org/10.1134/S2070048219060139>
18. H. Ezzin, B. Wang, Z. Qian, Propagation behavior of ultrasonic Love waves in functionally graded piezoelectric-piezomagnetic materials with exponential variation. *Mech. Mater.* **148**. 103492 (2020). <https://doi.org/10.1016/j.mechmat.2020.103492>
19. S. Pshenichnov, R. Ivanov, and M. Datcheva, Transient wave propagation in functionally graded viscoelastic structures. *Mathematics* **10**(23). Art. 4505 (2022). <https://doi.org/10.3390/math10234505>



LICENSE TO PUBLISH AGREEMENT FOR CONFERENCE PROCEEDINGS

This License to Publish must be signed and returned to the Proceedings Editor before the manuscript can be published. If you have questions about how to submit the form, please contact the AIP Publishing Conference Proceedings office (confproc@aip.org). For questions regarding the copyright terms and conditions of this License, please contact AIP Publishing's Office of Rights and Permissions, 1305 Walt Whitman Road, Suite 300, Melville, NY 11747-4300 USA; Phone 516-576-2268; Email: rights@aip.org.

Article Title ("Work"): Study of Non-Stationary Processes in Inhomogeneous Viscoelastic Materials
(Please indicate the final title of the Work. Any substantive changes made to the title after acceptance of the Work may require the completion of a new agreement.)

All Author(s): Sergey G. Pshenichnov, Ekaterina A. Korovaytseva, Anastasiya M. Razarenova
(Please list all the authors' names in order as they will appear in the Work. All listed authors must be fully deserving of authorship and no such authors should be omitted. For large groups of authors, attach a separate list to this form.)

Title of Conference: AMSMT 2025

Name(s) of Editor(s): Valentin L. Popov

All Copyright Owner(s), if not Author(s):

(Please list all copyright owner(s) by name. In the case of a Work Made for Hire, the employer(s) or commissioning party(ies) are the copyright owner(s). For large groups of copyright owners, attach a separate list to this form.)

Copyright Ownership and Grant of Rights

For the purposes of this License, the "Work" consists of all content within the article itself and made available as part of the article, including but not limited to the abstract, tables, figures, graphs, images, and multimedia files, as well as any subsequent errata. "Supplementary Material" consists of material that is associated with the article but linked to or accessed separately (available electronically only), including but not limited to data sets and any additional files.

This Agreement is an Exclusive License to Publish not a Transfer of Copyright. Copyright to the Work remains with the Author(s) or, in the case of a Work Made for Hire, with the Author(s)' employer(s). AIP Publishing LLC shall own and have the right to register in its name the copyright to the proceedings issue or any other collective work in which the Work is included. Any rights granted under this License are contingent upon acceptance of the Work for publication by AIP Publishing. If for any reason and at its own discretion AIP Publishing decides not to publish the Work, this License is considered void.

Each Copyright Owner hereby grants to AIP Publishing LLC the following irrevocable rights for the full term of United States and foreign copyrights (including any extensions):

1. The exclusive right and license to publish, reproduce, distribute, transmit, display, store, translate, edit, adapt, and create derivative works from the Work (in whole or in part) throughout the world in all formats and media whether now known or later developed, and the nonexclusive right and license to do the same with the Supplementary Material.
2. The right for AIP Publishing to freely transfer and/or sublicense any or all of the exclusive rights listed in #1 above. Sublicensing includes the right to authorize requests for reuse of the Work by third parties.
3. The right for AIP Publishing to take whatever steps it considers necessary to protect and enforce, at its own expense, the exclusive rights granted herein against third parties.

Author Rights and Permitted Uses

Subject to the rights herein granted to AIP Publishing, each Copyright Owner retains ownership of copyright and all other proprietary rights such as patent rights in the Work.

Each Copyright Owner retains the following nonexclusive rights to use the Work, without obtaining permission from AIP Publishing, in keeping with professional publication ethics and provided clear credit is given to its first publication in an AIP Publishing proceeding. Any reuse must include a full credit line acknowledging AIP Publishing's publication and a link to the Version of Record (VOR) on AIP Publishing's site.

Each Copyright Owner may:

1. Reprint portions of the Work (excerpts, figures, tables) in future works created by the Author, in keeping with professional publication ethics.
2. Post the Accepted Manuscript (AM) to their personal web page or their employer's web page immediately after acceptance by AIP Publishing.
3. Deposit the AM in an institutional or funder-designated repository immediately after acceptance by AIP Publishing.

4. Use the AM for posting within scientific collaboration networks (SCNs). For a detailed description of our policy on posting to SCNs, please see our Web Posting Guidelines (<https://publishing.aip.org/authors/web-posting-guidelines>).
5. Reprint the Version of Record (VOR) in print collections written by the Author, or in the Author's thesis or dissertation. It is understood and agreed that the thesis or dissertation may be made available electronically on the university's site or in its repository and that copies may be offered for sale on demand.
6. Reproduce copies of the VOR for courses taught by the Author or offered at the institution where the Author is employed, provided no fee is charged for access to the Work.
7. Use the VOR for internal training and noncommercial business purposes by the Author's employer.
8. Use the VOR in oral presentations made by the Author, such as at conferences, meetings, seminars, etc., provided those receiving copies are informed that they may not further copy or distribute the Work.
9. Distribute the VOR to colleagues for noncommercial scholarly use, provided those receiving copies are informed that they may not further copy or distribute the Work.
10. Post the VOR to their personal web page or their employer's web page 12 months after publication by AIP Publishing.
11. Deposit the VOR in an institutional or funder-designated repository 12 months after publication by AIP Publishing.
12. Update a prior posting with the VOR on a noncommercial server such as arXiv, 12 months after publication by AIP Publishing.

Author Warranties

Each Author and Copyright Owner represents and warrants to AIP Publishing the following:

1. The Work is the original independent creation of each Author and does not infringe any copyright or violate any other right of any third party.
2. The Work has not been previously published and is not being considered for publication elsewhere in any form, except as a preprint on a noncommercial server such as arXiv, or in a thesis or dissertation.
3. Written permission has been obtained for any material used from other sources and copies of the permission grants have been supplied to AIP Publishing to be included in the manuscript file.
4. All third-party material for which permission has been obtained has been properly credited within the manuscript.
5. In the event that the Author is subject to university open access policies or other institutional restrictions that conflict with any of the rights or provisions of this License, such Author has obtained the necessary waiver from his or her university or institution.

This License must be signed by the Author(s) and, in the case of a Work Made for Hire, also by the Copyright Owners. One Author/Copyright Owner may sign on behalf of all the contributors/owners only if they all have authorized the signing, approved of the License, and agreed to be bound by it. The signing Author and, in the case of a Work Made for Hire, the signing Copyright Owner warrants that he/she/it has full authority to enter into this License and to make the grants this License contains.

1. The Author must please sign here (except if an Author is a U.S. Government employee, then please sign under #3 below):

Author(s) Signature: [Signature] Print Name: Sergey Pshenichnov Date: 30.09.2025

2. The Copyright Owner (if different from the Author) must please sign here:

Name of Copyright Owner: _____ Authorized Signature and Title: _____ Date: _____

3. If an Author is a U.S. Government employee, such Author must please sign below. The signing Author certifies that the Work was written as part of his/her official duties and is therefore not eligible for copyright protection in the United States.

Name of U.S. Government Institution (e.g., Naval Research Laboratory, NIST): _____

Author Signature: _____ Print Name: _____ Date: _____

PLEASE NOTE: NATIONAL LABORATORIES THAT ARE SPONSORED BY U.S. GOVERNMENT AGENCIES BUT ARE INDEPENDENTLY RUN ARE NOT CONSIDERED GOVERNMENT INSTITUTIONS. (For example, Argonne, Brookhaven, Lawrence Livermore, Sandia, and others.) Authors at these types of institutions should sign under #1 or #2 above.

If the Work was authored under a U.S. Government contract, and the U.S. Government wishes to retain for itself and others acting on its behalf, a paid-up, nonexclusive, irrevocable, worldwide license in the Work to reproduce, prepare derivative works from, distribute copies to the public, perform publicly, and display publicly, by or on behalf of the Government, please check the box below and add the relevant Contract numbers.

☐ Contract #(s): _____ [11.18.1]

LICENSE TERMS DEFINED

Accepted Manuscript (AM): The final version of an author's manuscript that has been accepted for publication and incorporates all the editorial changes made to the manuscript after submission and peer review. The AM does not yet reflect any of the publisher's enhancements to the work such as copyediting, pagination, and other standard formatting.

arXiv: An electronic archive and distribution server for research article preprints in the fields of physics, mathematics, computer science, quantitative biology, quantitative finance, and statistics, which is owned and operated by Cornell University, <http://arxiv.org/>.

Commercial and noncommercial scholarly use: Noncommercial scholarly uses are those that further the research process for authors and researchers on an individual basis for their own personal purposes. They are author-to-author interactions meant for the exchange of ideas. Commercial uses fall outside the author-to-author exchange and include but are not limited to the copying or distribution of an article, either in hard copy form or electronically, for resale or licensing to a third party; posting of the AM or VOR of an article by a site or service where an access fee is charged or which is supported by commercial paid advertising or sponsorship; use by a for-profit entity for any type of promotional purpose. Commercial uses require the permission of AIP Publishing.

Embargo period: The period of time during which free access to the full text of an article is delayed.

Employer's web page: A web page on an employer's site that highlights the accomplishments and research interests of the company's employees, which usually includes their publications. (See also: Personal web page and Scholarly Collaboration Network).

Exclusive License to Publish: An exclusive license to publish is a written agreement in which the copyright owner gives the publisher exclusivity over certain inherent rights associated with the copyright in the work. Those rights include the right to reproduce the work, to distribute copies of the work, to perform and display the work publicly, and to authorize others to do the same. The publisher does not hold the copyright to the work, which continues to reside with the author. The terms of the AIP Publishing License to Publish encourage authors to make full use of their work and help them to comply with requirements imposed by employers, institutions, and funders.

Full Credit Line: AIP Publishing's preferred format for a credit line is as follows (you will need to insert the specific citation information in place of the capital letters): "Reproduced from [FULL CITATION], with the permission of AIP Publishing." A FULL CITATION would appear as: Journal abbreviation, volume number, article ID number or page number (year). For example: Appl. Phys. Lett. 107, 021102 (2015).

Institutional repository: A university or research institution's digital collection of articles that have been authored by its staff and which are usually made publicly accessible. As authors are encouraged and sometimes required to include their published articles in their institution's repository, the majority of publishers allow for deposit of the Accepted Manuscript for this purpose. AIP Publishing also allows for the VOR to be deposited 12 months after publication of the Work.

Journal editorial office: The contact point for authors concerning matters related to the publication of their manuscripts. Contact information for the journal editorial offices may be found on the journal websites under the "About" tab.

Linking to the Version of Record (VOR): To create a link to your article in an AIP Publishing journal or proceedings, you need to know the CrossRef digital object identifier (doi). You can find the doi on the article's abstract page. For instructions on linking, please refer to our Web Posting Guidelines at <https://publishing.aip.org/authors/web-posting-guidelines>.

National Laboratories: National laboratories are sponsored and funded by the U.S. Government but have independent nonprofit affiliations and employ private sector resources. These institutions are classified as Federally Funded Research and Development Centers (FFRDCs). Authors working at FFRDCs are not

considered U.S. Government employees for the purposes of copyright. The Master Government List of FFRDCs may be found at <http://www.nsl.gov/statistics/ffrdclist/>.

Personal web page: A web page that is hosted by the author or the author's institution and is dedicated to the author's personal research interests and publication history. An author's profile page on a social media site or scholarly collaboration network site is not considered a personal web page. (See also: Scholarly Collaboration Network; Employer's web page).

Peer X-Press: A web-based manuscript submission system by which authors submit their manuscripts to AIP Publishing for publication, communicate with the editorial offices, and track the status of their submissions. The Peer X-Press system provides a fully electronic means of completing the License to Publish. A hard copy of the Agreement will be supplied by the editorial office if the author is unable to complete the electronic version of the form. (Conference Proceedings authors will continue to submit their manuscripts and forms directly to the Conference Editors.)

Preprint: A version of an author's manuscript intended for publication but that has not been peer reviewed and does not reflect any editorial input or publisher enhancements.

Professional Publication Ethics: AIP Publishing provides information on what it expects from authors in its "Statement of ethics and responsibilities of authors submitting to AIP Publishing journals" (<http://publishing.aip.org/authors/ethics>). AIP Publishing is also a member of the Committee on Publication Ethics (COPE) (<http://publicationethics.org/>), which provides numerous resources and guidelines for authors, editors, and publishers with regard to ethical standards and accepted practices in scientific publishing.

Scholarly Collaboration Network (SCN): Professional networking sites that facilitate collaboration among researchers as well as the sharing of data, results, and publications. SCNs include sites such as Academia.edu, ResearchGate, and Mendeley, among others.

Supplementary Material: Related material that has been judged by peer review as being relevant to the understanding of the article but that may be too lengthy or of too limited interest for inclusion in the article itself. Supplementary Material may include data tables or sets, appendixes, movie or audio clips, or other multimedia files.

U.S. Government employees: Authors working at Government organizations who author works as part of their official duties and who are not able to license rights to the Work, since no copyright exists. Government works are in the public domain within the United States.

Version of Record (VOR): The final published version of the article as it appears in the printed journal/proceedings or on the Citation website. It incorporates all editorial input, is formatted in the publisher's standard style, and is usually viewed in PDF form.

Waiver: A request made to a university or institution to exempt an article from its open-access policy requirements. For example, a conflict will exist with any policy that requires the author to grant a nonexclusive license to the university or institution that enables it to license the Work to others. In all such cases, the Author must obtain a waiver, which shall be included in the manuscript file.

Work: The "Work" is considered all the material that comprises the article, including but not limited to the abstract, tables, figures, images, multimedia files that are directly embedded within the text, and the text itself. The Work does not include the Supplementary Material (see Supplementary Material above).

Work Made for Hire: Under copyright law, a work prepared by an employee within the scope of employment, or a work that has been specially ordered or commissioned for which the parties have agreed in writing to consider as a Work Made for Hire. The hiring party or employer is considered the author and owner of the copyright, not the person who creates the work.