

3rd International Conference Advanced Mechanics: Structure, Materials, Tribology

Application of Deep Learning and Physics-Informed Neural Networks to the Heat Transfer Problem within the Green-Naghdi Type II Theory

AIPCP25-CF-AMSMT2025-00022 | Article

PDF auto-generated using **ReView**



Application of Deep Learning and Physics-Informed Neural Networks to the Heat Transfer Problem within the Green–Naghdi Type II Theory

Gregory V. Fedotenkov^{1,2, a)} and Yana A. Vahterova^{1, b)}

¹*Moscow Aviation Institute (National Research University), 125993, Volokolamskoye shosse, 4, Moscow, Russia*

²*Institute of Mechanics of Lomonosov Moscow State University, 119192, Michurinsky prosp, 1, Moscow, Russia*

^{a)} Corresponding author: greghome@mail.ru
^{b)} yana-vahterova@mail.ru

Abstract. The paper considers the heat transfer problem within the framework of the Green–Naghdi type II theory, which accounts for the finite speed of thermal wave propagation. A comparison is made between the analytical solution, the numerical method based on a finite difference scheme, and the approach utilizing Physics-Informed Neural Networks (PINN). To validate the results, the parameters of duralumin are used in a dimensionless form. It is shown that the PINN method accurately reproduces the dynamics of thermal waves and demonstrates efficiency comparable to classical methods, while providing additional robustness to discretization errors. The results obtained confirm the prospects of applying PINN to solve both direct and inverse problems of hyperbolic heat conduction.

INTRODUCTION

The solution of hyperbolic heat conduction equations can be obtained using various methods. Analytical approaches, based on Fourier and Laplace transforms or Fourier series expansions, provide rigorous results but are applicable only to relatively simple geometries and boundary conditions. Numerical methods (finite-difference and finite-element schemes) make it possible to efficiently solve problems in complex domains and under arbitrary loading, but they require stability analysis, the coordination of time and space steps, as well as significant computational resources. In particular, methods based on second-order schemes in time and space have become widespread in solving dynamic heat conduction problems [5-6], [12-13], [23], [28-30].

In recent years, methods based on deep machine learning have been actively developed. Special attention has been drawn to the Physics-Informed Neural Networks (PINN) technology proposed in [20-22]. The main idea of PINN is the integration of physical laws into the neural network loss function, which makes it possible to obtain approximate solutions of partial differential equations without explicit discretization of the domain. This approach has proven effective for solving both direct and inverse problems, including heat conduction, elasticity, and acoustics [1-4], [7-11], [14-27]. Several studies have demonstrated that PINN is capable of reproducing complex dynamic processes and shows robustness to noisy data, which makes it a promising tool for engineering applications.

Of particular interest is the application of PINN to hyperbolic heat conduction models, in particular to the Green–Naghdi type II theory. Such formulations not only allow for a correct description of thermal wave propagation processes but also enable the solution of inverse problems related to the identification of material parameters or localization of heat sources.

In this paper, we consider a heat transfer problem within the framework of the Green–Naghdi type II theory with prescribed initial conditions and Dirichlet boundary conditions. Three solution approaches are compared: an analytical method (Fourier series expansion), a numerical method (second-order finite-difference scheme), and a neural network-based approach utilizing PINN. For verification purposes, a thermally conductive medium made of duralumin is considered, with its parameters given in dimensionless form. The main focus is placed on analyzing the accuracy, stability, and specific features of applying PINN to the modeling of thermal processes.

MATHEMATICAL FORMULATION OF THE PROBLEM

Let us consider the heat transfer problem in a homogeneous isotropic medium, described by the Green–Naghdi type II equation [3] (Fig. 1).




FIGURE 1. Illustration of the mathematical formulation of the problem.

In dimensional form, the governing equation can be written as:

$$\rho c_e \dot{\theta}^* = \tilde{\kappa}^* \square \theta^* + \rho q^{*(e)}, \quad (1)$$

where, ρ — is the material density, c_e — is the specific heat capacity at constant strain, T_0 — is the initial temperature, $\theta^*(x^*, t^*) = T^*(x^*, t^*) - T_0$ — is the temperature deviation from the initial value, $\tilde{\kappa}^*$ — is the thermal wave speed, $q^{*(e)}$ — is the volumetric heat source (the heat released per unit volume per unit time), and $\dot{q}^{*(e)}$ — is the rate of change of the heat source intensity with respect to time.

The Dirichlet boundary conditions are specified as:

$$\theta|_{x=0} = 0, \quad \theta|_{x=l} = 0, \quad (2)$$

and the initial conditions at $t^* = 0$:

$$\theta^*(x^*, 0) = T_0 \sin\left(\frac{\pi x^*}{l}\right), \quad \dot{\theta}^*(x^*, 0) = \frac{T_0 c_T}{l} \left(\frac{x^*}{l}\right) \left(\frac{x^*}{l} - 1\right), \quad (3)$$

For simplification of the analysis and for obtaining the canonical form of the governing equation, let us introduce dimensionless variables. As characteristic scales we choose the length l , the thermal wave propagation time l/c_T and the initial temperature T_0 . Then, the system of dimensionless variables takes the form:

$$x = \frac{x^*}{l}, \quad t = \frac{c_T t^*}{l}, \quad c_T^2 = \frac{\tilde{\kappa}}{\rho c_e}, \quad \theta = \frac{\theta^*}{T_0}, \quad q^{*(e)} = \frac{q^{*(e)} l}{c_e c_T T_0}. \quad (4)$$

where x, t — are the dimensionless spatial and temporal variables, θ — is the relative temperature deviation, c_T — is the thermal wave speed, and $q^{*(e)}$ — is the dimensionless external heat source.

Thus, the mathematical formulation of the problem in its dimensionless form is:

$$\ddot{\theta} - \square \theta = \dot{q}^{*(e)}, \quad x \in [0, 1], \quad t > 0, \quad (5)$$

with Dirichlet boundary conditions:

$$\theta|_{x=0} = \sin(\pi x), \quad \dot{\theta}|_{x=0} = x(x-1), \quad (6)$$

and initial conditions:

$$\theta|_{x=0} = 0, \quad \theta|_{x=l} = 0. \quad (7)$$

ANALYTICAL SOLUTION FOR THE GREEN–NAGHDI TYPE II HEAT CONDUCTION EQUATION

Let us consider the one-dimensional heat transfer problem within the framework of the Green–Naghdi type II theory in its dimensionless form (equations (5)–(7)). As an external heat source, we take the function:

$$q^{*(e)}(x, t) = 2xt, \quad (8)$$

which models a linearly increasing source in both space and time. In this case, the intensity of heat generation grows proportionally to the coordinate x and time t , which corresponds to a non-stationary process of localized heat release.

Then its time derivative, which enters the governing equation (5), takes the form:

$$\dot{q}(x, t) = 2x. \quad (9)$$

To solve the problem (5) – (7) we apply the method of separation of variables with the expansion of the desired function in terms of the eigenfunctions of the Sturm–Liouville boundary value problem:

$$\vartheta(x, t) = \sum_{n=1}^{\infty} a_n(t) \sin(\pi n x). \quad (10)$$

Substituting this representation into the governing equation leads to the following system for the coefficients:

$$a_n''(t) + (\pi n)^2 a_n(t) = F_n, \quad F_n = 4 \int_0^1 x \sin(\pi n x) dx = \frac{4(-1)^{n+1}}{\pi n}, \quad (11)$$

where F_n are the Fourier coefficients of the source term.

At $t = 0$ the temperature distribution in the rod is prescribed as a sinusoidal function (6). This condition corresponds to a perturbation in which the maximum temperature deviation from the initial value occurs at the middle of the domain, while at the ends $x = 0$ and $x = 1$ the temperature vanishes due to the imposed Dirichlet boundary conditions. In addition, (6) specifies the distribution of the initial rate of temperature change. This quadratic function is negative in the interval $0 \leq x \leq 1$ and vanishes at the boundaries, which is consistent with the end-fixing conditions. Thus, at time $t = 0$ the temperature is not only distributed according to a sinusoidal law, but also begins to change at a given rate, which reflects the dynamic nature of the problem statement.

When moving to the coefficients of expansion in eigenfunctions, the system of initial conditions takes the form

$$a_n(0) = \delta_{n1}, \quad a_n'(0) = \begin{cases} 0, & n - \text{even number}, \\ -\frac{8}{\pi^3 n^3}, & n - \text{odd number}, \end{cases} \quad (12)$$

where δ_{n1} — is the Kronecker delta, equal to 1 for $n = 1$ and 0 otherwise.

The solution for each coefficient a_n from (11) is:

$$a_n(t) = \left(\delta_{n1} - \frac{4(-1)^{n+1}}{\pi^3 n^3} \right) \cos(\pi n t) + \frac{4((-1)^n - 1)}{\pi^4 n^4} \sin(\pi n t) + \frac{4(-1)^{n+1}}{\pi^3 n^3}. \quad (13)$$

Thus, the general solution of the problem is represented as a Fourier series:

$$\vartheta(x, t) = \sum_{n=1}^{\infty} \left[\left(\delta_{n1} - \frac{4(-1)^{n+1}}{\pi^3 n^3} \right) \cos(\pi n t) + \frac{4((-1)^n - 1)}{\pi^4 n^4} \sin(\pi n t) + \frac{4(-1)^{n+1}}{\pi^3 n^3} \right] \sin(\pi n x). \quad (14)$$

NUMERICAL SOLUTION FOR THE GREEN–NAGHDI TYPE II HEAT CONDUCTION EQUATION

For the numerical solution of the one-dimensional heat transfer problem within the framework of the Green–Naghdi type II theory (equations (5)–(7)) we introduce uniform grids in the spatial coordinate and time:

$$x_j = j \Delta x, \quad j = 0, \dots, M, \quad t^n = n \Delta t, \quad n = 0, 1, \dots \quad (15)$$

and seek approximations $u_j^n \approx \vartheta(x_j, t^n)$. The second derivative with respect to space is approximated by a central difference of second order:

$$\vartheta_{xx}(x_j, t^n) \approx \frac{u_{j+1}^n - 2u_j^n + u_{j-1}^n}{\Delta x^2}. \quad (16)$$

Applying central differences in time leads to an explicit scheme of second-order accuracy:

$$u_j^{n+1} = 2u_j^n - u_j^{n-1} + \lambda^2 (u_{j+1}^n - 2u_j^n + u_{j-1}^n) + (\Delta t)^2 f_j, \quad j = 1, \dots, M-1, \quad (17)$$

where $\lambda = \Delta t / \Delta x$, and $f_j = 2x_j$. The Dirichlet boundary conditions are imposed explicitly:

$$u_0^n = 0, \quad u_M^n = 0, \quad n \geq 0. \quad (18)$$

To construct the first time layer, a Taylor expansion is used with account of the governing equation:

$$u_j^1 = u_j^0 + \Delta t, v_j^0 + \frac{1}{2}(\Delta t)^2 \left((\vartheta_{xx})_j^0 + f_j \right), \quad (19)$$

where $u_j^0 = \sin(\pi x_j)$, $v_j^0 = x_j(x_j - 1)$.

A Fourier analysis for the homogeneous equation shows that the scheme is stable under the Courant–Friedrichs–Lowy condition:

$$\lambda = \frac{\Delta t}{\Delta x} \leq 1. \quad (20)$$

Thus, the time step is selected from condition (20). The scheme has second-order accuracy in space and time.

To control accuracy, an analytical solution in the form of a Fourier series of eigenfunctions of the Dirichlet problem was used. Comparison of the numerical results with a truncated series (400 members) showed that the relative error in the norm of L_2 on the time interval $0 \leq t \leq 1.2$ does not exceed 10^{-6} . This confirms the correctness of the implementation and the second-order convergence with respect to the grid steps.

SOLUTION OF THE GREEN–NAGHDI TYPE II HEAT CONDUCTION EQUATION USING A NEURAL NETWORK

PINN approximation with exact enforcement of boundary and initial conditions

To satisfy the boundary and initial conditions exactly, a special representation of the solution is used, allowing them to be embedded directly into the model structure:

$$\vartheta(x, t) = g(x) + th(x) + t^2 S(x) N_\phi(x, t), \quad (21)$$

where $S(x) = x(1-x)$ vanishes at the boundary $x = 0, 1$, and $N_\phi : \mathbb{D}^2 \rightarrow \mathbb{D}$ — is the output of a fully connected neural network with parameters ϕ .

Verification of conditions:

Boundary conditions — $\vartheta(0, t) = g(0) + th(0) + t^2 S(0) N_\phi(0, t) = 0$ and similarly $\vartheta(1, t) = 0$.

Initial conditions — $\vartheta(x, 0) = g(x)$, because $t = 0$ vanishes the terms with $h(x)$ и N_ϕ ;

further $\vartheta_t(x, t) = h(x) + 2tS(x)N_\phi + t^2 S(x)\partial_t N_\phi$, where $\vartheta_t(x, 0) = h(x)$.

Thus, the loss function contains no additional penalty terms for boundary and initial conditions, and the training of the neural network is focused exclusively on the dynamics of the equation.

The approximating model is a fully connected neural network (Multi-Layer Perceptron, MLP). The inputs are the spatial and temporal coordinates $(x, t) \in \mathbb{D}^2$, which ensures joint modeling of the solution's dependence on both spatial and temporal variables. The output of the network is a scalar value $N_\phi(x, t)$, representing a part of the solution in the constructive form.

Network architecture and initialization

The network architecture includes five hidden layers with 128 neurons each, with the \tanh activation function. This choice is motivated by its properties: smoothness, boundedness, and the ability to adequately reproduce both low-frequency and moderately oscillatory components of the solution. Such a combination of depth and width provides sufficient approximation capacity while maintaining training stability.

Weights are initialized according to the Xavier (Glorot) scheme, which ensures a balanced scale of the initial parameters and promotes more stable convergence of the optimization process. For problems with a more complex spectral content (e.g., dominated by high-frequency components), alternative activation functions such as sinusoidal ones (SIREN), as well as an adapted weight initialization strategy, are preferable. This approach improves the ability of the model to reproduce oscillatory solution behavior, which is typical for hyperbolic-type problems.

Loss function and automatic differentiation

The training of the neural network model is based on minimizing the residual of the Green–Naghdi type II heat conduction equation [4]. The only functional used is the mean squared residual of the governing equation at the collocation points:

$$L_{\text{PDE}}(\phi) = \frac{1}{N_c} \sum_{i=1}^{N_c} \left[\vartheta_u(x_i, t_i; \phi) - \vartheta_{xx}(x_i, t_i; \phi) - \dot{q}^{(e)}(x_i, t_i) \right]^2, \quad (22)$$

where $\{(x_i, t_i)\}_{i=1}^{N_c}$ — is a sample of points from the region $x \in (0, 1), t \in (0, T]$, and the heat source has the form $\dot{q}^{(e)}(x, t) = 2x$.

The derivatives $\vartheta_t, \vartheta_u, \vartheta_x, \vartheta_{xx}$, are computed using automatic differentiation. In particular, in PyTorch this is implemented by successive calls of the *autograd.grad* function, which allows obtaining derivatives of any order with respect to spatial and temporal variables without explicit construction of finite-difference approximations. This significantly simplifies implementation and provides machine precision in computing gradients, which is necessary for training the network.

Special attention was paid to the stability of second derivative computations, since they are most sensitive to approximation errors and numerical artifacts. To improve accuracy, derivatives were computed in standard floating-point precision (float32), even when mixed precision (AMP) was used for acceleration.

It should be noted that mixed precision (AMP) can accelerate training on GPUs due to computations in float16/bfloat16 format. However, when calculating second derivatives, loss of accuracy and degradation of convergence are possible. Therefore, in the base configuration, AMP was disabled, and when enabled, additional quality control is required (e.g., comparison with reference values or fallback to float32 for derivative computation).

Thus, the loss function is formed solely by the residual of the equation, while the boundary and initial conditions are automatically satisfied through the special solution representation. This allows the optimization of network parameters to focus on approximating the equation dynamics, ensuring high accuracy and stability of training.

Strategy for generating collocation points

A key element of the PINN method is the generation of the set of collocation points $\{(x_i, t_i)\}_{i=1}^{N_c}$, at which the residual of the equation is evaluated. In the basic configuration, the points are uniformly sampled over the rectangular domain $(0, 1) \times (0, T]$. This approach ensures coverage of the entire integration domain and uniform distribution of errors in space and time.

To improve training stability, resampling is applied: at each training iteration, a new set of collocation points is generated. This makes it possible to:

- reduce correlation between successive batches,
- avoid “fitting” the solution to a fixed grid,
- increase robustness against trapping in local minima of the loss function.

In some cases, it is advisable to use more advanced point generation schemes, such as Latin Hypercube Sampling (LHS) or Sobol sequences. These methods provide better filling of the domain and allow for more uniform coverage of the variable space than simple uniform sampling.

When solving problems with pronounced solution fronts, characteristic of hyperbolic equations, local point refinement is useful. In particular, it is recommended to increase the density of collocation points:

- near the initial moment of time $t \approx 0$, where the solution is most sensitive to the initial perturbation,
- near the boundary points $x = 0, 1$, where strict boundary conditions are imposed.

Thus, an adaptive sampling strategy significantly improves the agreement of the PINN solution with the analytical reference and accelerates training convergence.

The training of the neural network model is carried out on a graphics processor and is organized in two phases:

– **Phase 1 (Adam).** At the first stage, the stochastic gradient optimizer Adam [24] is used with an initial learning rate $lr = 10^{-3}$. To control convergence, an adaptive scheduler ReduceLROnPlateau is applied, which automatically decreases the learning rate in the absence of noticeable progress. A minimum of 2000 epochs is allocated for the “warm-up” of the model, allowing the network to leave the random initialization state and to form a correct approximation.

– **Phase 2 (L-BFGS).** At the second stage, fine-tuning of the parameters is performed using the quasi-Newton method L-BFGS with line search according to the “strong Wolfe” condition. This method effectively minimizes quadratic functionals and allows achieving high solution accuracy near the optimum.

The transition between training phases is implemented using a combination of criteria:

- *early transition*: if the equation residual becomes smaller than the threshold $L_{PDE} < 10^{-4}$;
- *plateau detection*: if the improvement of the error is less than 10^{-5} for 800 consecutive epochs;
- *learning rate condition*: if the learning rate decreases below a prescribed threshold $lr < 5 \cdot 10^{-5}$.

Such a hybrid strategy adapts the training process to the dynamics of the specific problem: Adam ensures rapid descent in the initial stages, while L-BFGS refines the solution to high accuracy. As a result, computational resources are saved and a stable reduction of the residual is achieved while maintaining accuracy compared to analytical and numerical solutions.

CALCULATION RESULTS

To verify the correctness of the proposed PINN scheme, computational experiments were carried out for the one-dimensional heat transfer problem within the framework of the Green–Naghdi type II theory. As reference solutions, the following were used:

1. the analytical solution represented as a Fourier series in trigonometric functions (14);
2. an independent numerical solution (17), obtained by a finite-difference wave-type scheme.

The PINN model was trained on a graphics processor in a two-phase mode: in the first stage the Adam $lr = 10^{-3}$ algorithm with ReduceLROnPlateau learning rate reduction was applied, and in the second stage fine-tuning was performed using the L-BFGS method (Fig. 2).




FIGURE 2. Loss function history during PINN training: Adam phase and fine-tuning with L-BFGS.

The minimization of the loss function was stable, with a monotonic decrease of the residual and achievement of the threshold value $L_{PDE} < 10^{-4}$.

Comparison of the solutions showed a high degree of agreement between the PINN results and the analytical and finite-difference solutions. For a quantitative assessment of accuracy, global relative errors in the L_2 – norm were calculated (Fig. 3):

$$\varepsilon_{FD/Analytic} \approx O(10^{-3}), \varepsilon_{PINN/Analytic} \approx O(10^{-4}), \varepsilon_{PINN/FD} \approx O(10^{-3}) \quad (23)$$




FIGURE 3. Relative L_2 - error between the analytical solution, finite-difference approximation, and the PINN model.

The obtained values demonstrate comparable accuracy of PINN and the classical finite-difference approximation, while PINN does not require explicit discretization of the domain.

The graphical results are presented as solution cross-sections with respect to spatial and temporal variables $\theta(x_0, t)$ at fixed values of x_0 and $\theta(x, t_0)$ at fixed instants of time t_0 (Fig. 4). In these graphs, the solutions obtained by different methods practically coincide.




FIGURE 4. Temperature field $\theta(x_0, t)$ for fixed x_0 and $\theta(x, t_0)$ for fixed t_0 .

Additionally, three-dimensional surfaces $\theta(x, t)$ were constructed for both the PINN and analytical solutions (Fig. 5). The analysis of these visualizations confirms the correct reproduction by PINN of the wave nature of the Green–Naghdi type II model, including the finite propagation speed of thermal disturbances.




FIGURE 5. Three-dimensional surfaces of the temperature field $\theta(x, t)$, obtained using PINN and the analytical solution.

Thus, the performed calculations showed that the proposed approach provides high approximation accuracy and can be considered as an alternative tool to classical numerical schemes for solving hyperbolic heat conduction problems.

CONCLUSION

In this work, the one-dimensional heat transfer problem within the framework of the Green–Naghdi type II theory was considered using Physics-Informed Neural Networks (PINN). A neural network approximation of the solution was constructed based on a special representation that ensures exact satisfaction of the boundary and initial conditions. The mean squared residual of the governing equation, computed at collocation points using automatic differentiation, was employed as the loss function.

The conducted computational experiments showed that the proposed approach makes it possible to obtain a solution comparable in accuracy to the analytical Fourier series and the independent finite-difference method. The relative error in the L_2 – norm for PINN does not exceed 10^{-3} , which indicates the high accuracy of the method. Graphical analysis (cross-sections with respect to spatial and temporal variables, three-dimensional surfaces of the temperature field) confirmed the correct reproduction of the wave nature of the Green–Naghdi type II model.

The proposed two-phase training scheme with optimization Adam → L-BFGS and resampling of collocation points ensures stable convergence and effective reduction of the residual. Thus, PINN can be considered as a universal tool for solving hyperbolic heat conduction problems, combining the accuracy of analytical methods with the flexibility of numerical schemes.

Promising directions for further research include extending the proposed approach to multidimensional problems and more complex geometries, as well as the use of sinusoidal activations (SIREN), which improve the reproduction of high-frequency components of the solution. In addition, the method can be effectively applied to inverse problems of material parameter identification and heat source localization. Of particular interest is the analysis of the robustness of PINN to noisy data and the development of regularization schemes aimed at improving the reliability and accuracy of the approximation.

ACKNOWLEDGMENTS

The research was supported by Russian Science Foundation grant No. 23-19-00680, <https://rscf.ru/project/23-19-00680/>.

REFERENCES

1. A. Ivanova, N. Kharin, T. Baltina and O. Sachenkov, "Muscle tone control system based on LIF model neural network," VIII International Conference on Information Technology and Nanotechnology (ITNT), Samara, Russian Federation, (2022), pp. 1-4, doi: 10.1109/ITNT55410.2022.9848650

2. A. Ivanova, N. Kharin, V. Smirnova, E. Tufanova and O.A. Sachenkov, (2022). Stabilization of a pendulum on an elastic foundation using a multilayer perceptron. *Journal of Physics: Conference Series*. 2308. 012005. 10.1088/1742-6596/2308/1/012005
3. A. V. Lykov, Theory of thermal conductivity. – (1967). [in Russian]
4. A.N. Kolmogorov, *Dokl. AN SSSR* **114** (5), 953–956 (1957) URL: <https://www.mathnet.ru/rus/dan22050>.
5. A.O. Serdyuk, G.V. Fedotenkov, Unsteady bending function for an unlimited anisotropic plate // *Vestnik Samarskogo Gosudarstvennogo Tekhnicheskogo Universiteta, Seriya Fiziko-Matematicheskie Nauki*, Vol. **25**, No. 1, – pp. 111-126, (2021). <https://doi.org/10.14498/vsgtu1793>
6. A.V. Zemskov, D.V. Tarlakovskii, G.V. Fedotenkov, “One-dimensional unsteady problems of thermoelasticity: Textbook”, Moscow, Izd-vo MAI, (2023), 96 p. [in Russian]
7. C. Irrgang, N. Boers, M. Sonnewald, et al., “Towards neural Earth system modelling by integrating artificial intelligence in Earth system science”, *Nat. Mach. Intelligence* **3** (8), 667–674 (2021). <https://doi.org/10.1038/s42256-021-00374-3>
8. D. Yarotsky, “Error bounds for approximations with deep relu networks”, *Neural Netw.* **94**, 103–114 (2017), DOI: 10.1016/j.neunet.2017.07.002.
9. D. Zhang, L. Lu, L. Guo, et al. “Quantifying total uncertainty in physics-informed neural networks for solving forward and inverse stochastic problems”, *J. Comput. Phys.* **397**, 108850 (2019), DOI: 10.1016/j.jcp.2019.07.048.
10. G. E. Karniadakis, I. G. Kevrekidis, L. Lu, et al. “Physics-informed machine learning”, *Nature Reviews Phys.* **3** (6), 422–440 (2021), URL: www.nature.com/articles/s42254-021-00314-5. DOI: 10.1038/s42254-021-00314-5.
11. G. Pang, L. Lu, G. E. Karniadakis, “fPINNs: Fractional Physics-Informed Neural Networks”, *SIAM J. Sci. Comput.* **41** (4), 2603–2626 (2019), DOI: 10.1137/18M1229845.
12. G. V. Fedotenkov, D. V. Tarlakovsky, Y. A. Vahterova, “Identification of Non-stationary Load Upon Timoshenko Beam”, *Lobachevskii Journal of Mathematics* **40** (4), 439-447 (2019), DOI 10.1134/S1995080219040061.
13. G. V. Fedotenkov, Y. A. Vahterova, D. I. Makarevskii, T. Q. Thang, “The inverse non-stationary problem of identification of defects in an elastic rod”, *INCAS Bulletin* **13** (Special Issue), 57-66 (2021), DOI 10.13111/2066-8201.2021.13.S.6.
14. I. Lagaris, A. Likas, D. Fotiadis, “Artificial neural networks for solving ordinary and partial differential equations”, *IEEE Trans. Neural Networks* **9** (5), 987–1000 (1998). <https://doi.org/10.1109/72.712178>
15. J. Berg, K. Nyström, “A unified deep artificial neural network approach to partial differential equations in complex geometries”, *Neurocomputing* **317**, 28–41 (2018). <https://doi.org/10.1016/j.neucom.2018.06.056>
16. J. Braun, M. Griebel, “On a constructive proof of Kolmogorov’s superposition theorem”, *Constructive Approximation journal* **30** (2009), DOI: 10.1007/s00365-009-9054-2.
17. J. Blechschmidt, O.G. Ernst, “Three ways to solve partial differential equations with neural networks – A review”, *GAMM-Mitteilungen* **44** (2), e202100,006 (2021). DOI: 10.1002/gamm.202100006.
18. L. Lu, R. Pestourie, W. Yao, et al., “Physics-informed neural networks with hard constraints for inverse design”, *SIAM J. Sci. Comput.* **43** (6), 1105–1132 (2021), DOI: 10.1137/21M1397908.
19. L. Yang, D. Zhang, G. E. Karniadakis, “Physics-informed generative adversarial networks for stochastic differential equations”, *SIAM J. Sci. Comput.* **42** (1), 292–317 (2020), DOI: 10.1137/18M1225409.
20. M. Raissi, P. Perdikaris, G. E. Karniadakis, “Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations”, *J. Comput. Phys.* **378**, 686–707 (2019), DOI: 10.1016/j.jcp.2018.10.045.
21. M. Raissi, P. Perdikaris, G. E. Karniadakis, “Physics Informed Deep Learning (Part I): Data-driven Solutions of Nonlinear Partial Differential Equations”, arXiv: 1711.10561, (2017), URL: <https://arxiv.org/abs/1711.10561v1>, DOI :10.48550/arXiv.1711.10561
22. M. Raissi, P. Perdikaris, G. E. Karniadakis, “Physics Informed Deep Learning (Part II): Data-driven Discovery of Nonlinear Partial Differential Equations”, arXiv: 1711.10566, (2017), URL: <https://arxiv.org/abs/1711.10566>, DOI: 10.48550/arXiv.1711.10566.
23. N. A. Lokteva, D. O. Serdyuk and P.D. Skopintsev, Non-stationary influence function for an unbounded anisotropic Kirchoff–Love shell // *Journal of Applied Engineering Science*, Vol. **18**, No. 4, pp. 737-744. (2020). <https://doi.org/10.5937/jaes0-28205>
24. P. Kingma Diederik, Ba. Jimmy “Adam: A Method for Stochastic Optimization”, arXiv: 1412.6980, (2017). <https://arxiv.org/abs/1412.6980>, DOI: 10.48550/arXiv.1412.6980.

25. S. Cai, Z. Mao, Z. Wang, et al., “Physics-informed neural networks (PINNs) for fluid mechanics: a review”, *Acta. Mech. Sin.* **37** (12), 1727–1738 (2021). DOI: 10.1007/s10409-021-01148-1.
26. S. Hajkin Nejronnye seti: polnyj kurs 2-e izdanie, Moscow, Izdatel'skij dom «Vil'yams», (2006), 1104 p.[in Russian]
27. S. Kollmannsberger, D. D'Angella, M. Jokeit, et al., “Physics-Informed Neural Networks”, in: Kollmannsberger, S., D'Angella, D., Jokeit, M., et al. (eds.). Deep Learning in Computational Mechanics. Studies in Computational Intelligence **977**, 55–84 (2021). DOI: 10.1007/978-3-030-76587-3_5.
28. V. A. Vestyak, A. V. Zemskov, Elektronnij zhurnal “Trudy MAI” **53**, (2012), available at: <https://trudymai.ru/published.php?ID=29732>. [in Russian]
29. V.I. Goncharenko and V.S. Oleshko, The Use of Artificial Neural Networks in Nondestructive Testing of Aircraft Parts // Russian Aeronautics, Volume **67**. No. 3, pp. 497-503,(2024). DOI: 10.3103/S1068799824030048.
30. Y.A. Vahterova, G. V. Fedotenkov, “The inverse problem of recovering an unsteady linear load for an elastic rod of finite length”, *Journal of Applied Engineering Science* **18** (4), 687-692 (2020), DOI10.5937/jaes0-28073.



LICENSE TO PUBLISH AGREEMENT FOR CONFERENCE PROCEEDINGS

This License to Publish must be signed and returned to the Proceedings Editor before the manuscript can be published. If you have questions about how to submit the form, please contact the AIP Publishing Conference Proceedings office (confproc@aip.org). For questions regarding the copyright terms and conditions of this License, please contact AIP Publishing's Office of Rights and Permissions, 1305 Walt Whitman Road, Suite 300, Melville, NY 11747-4300 USA; Phone 516-576-2268; Email: rights@aip.org.

Article Title ("Work"): Application of Deep Learning and Physics-Informed Neural Networks to the Heat Transfer Problem within the Green-Naghdi Type II Theory

(Please indicate the final title of the Work. Any substantive changes made to the title after acceptance of the Work may require the completion of a new agreement.)

All Author(s): Gregory V. Fedotenkov, Yana A. Vahterova

(Please list **all** the authors' names in order as they will appear in the Work. All listed authors must be fully deserving of authorship and no such authors should be omitted. For large groups of authors, attach a separate list to this form.)

Title of Conference: 3rd International Conference Advanced Mechanics: Structure, Materials, Tribology

Name(s) of Editor(s) Valentin L. Popov

All Copyright Owner(s), if not Author(s):

(Please list **all** copyright owner(s) by name. In the case of a Work Made for Hire, the employer(s) or commissioning party(ies) are the copyright owner(s). For large groups of copyright owners, attach a separate list to this form.)

Copyright Ownership and Grant of Rights

For the purposes of this License, the "Work" consists of all content within the article itself and made available as part of the article, including but not limited to the abstract, tables, figures, graphs, images, and multimedia files, as well as any subsequent errata. "Supplementary Material" consists of material that is associated with the article but linked to or accessed separately (available electronically only), including but not limited to data sets and any additional files.

This Agreement is an Exclusive License to Publish not a Transfer of Copyright. Copyright to the Work remains with the Author(s) or, in the case of a Work Made for Hire, with the Author(s)' employer(s). AIP Publishing LLC shall own and have the right to register in its name the copyright to the proceedings issue or any other collective work in which the Work is included. Any rights granted under this License are contingent upon acceptance of the Work for publication by AIP Publishing. If for any reason and at its own discretion AIP Publishing decides not to publish the Work, this License is considered void.

Each Copyright Owner hereby grants to AIP Publishing LLC the following irrevocable rights for the full term of United States and foreign copyrights (including any extensions):

1. The exclusive right and license to publish, reproduce, distribute, transmit, display, store, translate, edit, adapt, and create derivative works from the Work (in whole or in part) throughout the world in all formats and media whether now known or later developed, and the nonexclusive right and license to do the same with the Supplementary Material.
2. The right for AIP Publishing to freely transfer and/or sublicense any or all of the exclusive rights listed in #1 above. Sublicensing includes the right to authorize requests for reuse of the Work by third parties.
3. The right for AIP Publishing to take whatever steps it considers necessary to protect and enforce, at its own expense, the exclusive rights granted herein against third parties.

Author Rights and Permitted Uses

Subject to the rights herein granted to AIP Publishing, each Copyright Owner retains ownership of copyright and all other proprietary rights such as patent rights in the Work.

Each Copyright Owner retains the following nonexclusive rights to use the Work, without obtaining permission from AIP Publishing, in keeping with professional publication ethics and provided clear credit is given to its first publication in an AIP Publishing proceeding. Any reuse must include a full credit line acknowledging AIP Publishing's publication and a link to the Version of Record (VOR) on AIP Publishing's site.

Each Copyright Owner may:

1. Reprint portions of the Work (excerpts, figures, tables) in future works created by the Author, in keeping with professional publication ethics.
2. Post the Accepted Manuscript (AM) to their personal web page or their employer's web page immediately after acceptance by AIP Publishing.
3. Deposit the AM in an institutional or funder-designated repository immediately after acceptance by AIP Publishing.

4. Use the AM for posting within scientific collaboration networks (SCNs). For a detailed description of our policy on posting to SCNs, please see our Web Posting Guidelines (<https://publishing.aip.org/authors/web-posting-guidelines>).
5. Reprint the Version of Record (VOR) in print collections written by the Author, or in the Author's thesis or dissertation. It is understood and agreed that the thesis or dissertation may be made available electronically on the university's site or in its repository and that copies may be offered for sale on demand.
6. Reproduce copies of the VOR for courses taught by the Author or offered at the institution where the Author is employed, provided no fee is charged for access to the Work.
7. Use the VOR for internal training and noncommercial business purposes by the Author's employer.
8. Use the VOR in oral presentations made by the Author, such as at conferences, meetings, seminars, etc., provided those receiving copies are informed that they may not further copy or distribute the Work.
9. Distribute the VOR to colleagues for noncommercial scholarly use, provided those receiving copies are informed that they may not further copy or distribute the Work.
10. Post the VOR to their personal web page or their employer's web page 12 months after publication by AIP Publishing.
11. Deposit the VOR in an institutional or funder-designated repository 12 months after publication by AIP Publishing.
12. Update a prior posting with the VOR on a noncommercial server such as arXiv, 12 months after publication by AIP Publishing.

Author Warranties

Each Author and Copyright Owner represents and warrants to AIP Publishing the following:

1. The Work is the original independent creation of each Author and does not infringe any copyright or violate any other right of any third party.
2. The Work has not been previously published and is not being considered for publication elsewhere in any form, except as a preprint on a noncommercial server such as arXiv, or in a thesis or dissertation.
3. Written permission has been obtained for any material used from other sources and copies of the permission grants have been supplied to AIP Publishing to be included in the manuscript file.
4. All third-party material for which permission has been obtained has been properly credited within the manuscript.
5. In the event that the Author is subject to university open access policies or other institutional restrictions that conflict with any of the rights or provisions of this License, such Author has obtained the necessary waiver from his or her university or institution.

This License must be signed by the Author(s) and, in the case of a Work Made for Hire, also by the Copyright Owners. One Author/Copyright Owner may sign on behalf of all the contributors/owners only if they all have authorized the signing, approval of the License, and agreed to be bound by it. The signing Author and, in the case of a Work Made for Hire, the signing Copyright Owner warrants that he/she/it has full authority to enter into this License and to make the grants this License contains.

1. The Author must please sign here (except if an Author is a U.S. Government employee, then please sign under #3 below):

 Gregory V. Fedotenkov October 01, 2025
Author(s) Signature Print Name Date

2. The Copyright Owner (if different from the Author) must please sign here:

Name of Copyright Owner Authorized Signature and Title Date

3. If an Author is a U.S. Government employee, such Author must please sign below. The signing Author certifies that the Work was written as part of his/her official duties and is therefore not eligible for copyright protection in the United States.

Name of U.S. Government Institution (e.g., Naval Research Laboratory, NIST)

Author Signature Print Name Date

PLEASE NOTE: NATIONAL LABORATORIES THAT ARE SPONSORED BY U.S. GOVERNMENT AGENCIES BUT ARE INDEPENDENTLY RUN ARE NOT CONSIDERED GOVERNMENT INSTITUTIONS. (For example, Argonne, Brookhaven, Lawrence Livermore, Sandia, and others.) Authors at these types of institutions should sign under #1 or #2 above.

If the Work was authored under a U.S. Government contract, and the U.S. Government wishes to retain for itself and others acting on its behalf, a paid-up, nonexclusive, irrevocable, worldwide license in the Work to reproduce, prepare derivative works from, distribute copies to the public, perform publicly, and display publicly, by or on behalf of the Government, please check the box below and add the relevant Contract numbers.

Contract #(s) _____ [1.16.1]

LICENSE TERMS DEFINED

Accepted Manuscript (AM): The final version of an author's manuscript that has been accepted for publication and incorporates all the editorial changes made to the manuscript after submission and peer review. The AM does not yet reflect any of the publisher's enhancements to the work such as copyediting, pagination, and other standard formatting.

arXiv: An electronic archive and distribution server for research article preprints in the fields of physics, mathematics, computer science, quantitative biology, quantitative finance, and statistics, which is owned and operated by Cornell University, <http://arxiv.org>.

Commercial and noncommercial scholarly use: *Noncommercial* scholarly uses are those that further the research process for authors and researchers on an individual basis for their own personal purposes. They are author-to-author interactions meant for the exchange of ideas. *Commercial* uses fall outside the author-to-author exchange and include but are not limited to the copying or distribution of an article, either in hard copy form or electronically, for resale or licensing to a third party; posting of the AM or VOR of an article by a site or service where an access fee is charged or which is supported by commercial paid advertising or sponsorship; use by a for-profit entity for any type of promotional purpose. Commercial uses require the permission of AIP Publishing.

Embargo period: The period of time during which free access to the full text of an article is delayed.

Employer's web page: A web page on an employer's site that highlights the accomplishments and research interests of the company's employees, which usually includes their publications. (See also: Personal web page and Scholarly Collaboration Network).

Exclusive License to Publish: An exclusive license to publish is a written agreement in which the copyright owner gives the publisher exclusivity over certain inherent rights associated with the copyright in the work. Those rights include the right to reproduce the work, to distribute copies of the work, to perform and display the work publicly, and to authorize others to do the same. The publisher does not hold the copyright to the work, which continues to reside with the author. The terms of the AIP Publishing License to Publish encourage authors to make full use of their work and help them to comply with requirements imposed by employers, institutions, and funders.

Full Credit Line: AIP Publishing's preferred format for a credit line is as follows (you will need to insert the specific citation information in place of the capital letters): "Reproduced from [FULL CITATION], with the permission of AIP Publishing." A FULL CITATION would appear as: Journal abbreviation, volume number, article ID number or page number (year). For example: Appl. Phys. Lett. 107, 021102 (2015).

Institutional repository: A university or research institution's digital collection of articles that have been authored by its staff and which are usually made publicly accessible. As authors are encouraged and sometimes required to include their published articles in their institution's repository, the majority of publishers allow for deposit of the Accepted Manuscript for this purpose. AIP Publishing also allows for the VOR to be deposited 12 months after publication of the Work.

Journal editorial office: The contact point for authors concerning matters related to the publication of their manuscripts. Contact information for the journal editorial offices may be found on the journal websites under the "About" tab.

Linking to the Version of Record (VOR): To create a link to your article in an AIP Publishing journal or proceedings, you need to know the CrossRef digital object identifier (doi). You can find the doi on the article's abstract page. For instructions on linking, please refer to our Web Posting Guidelines at <https://publishing.aip.org/authors/web-posting-guidelines>.

National Laboratories: National laboratories are sponsored and funded by the U.S. Government but have independent nonprofit affiliations and employ private sector resources. These institutions are classified as Federally Funded Research and Development Centers (FFRDCs). Authors working at FFRDCs are not

considered U.S. Government employees for the purposes of copyright. The Master Government List of FFRDCs may be found at <http://www.nsf.gov/statistics/ffrclist/>.

Personal web page: A web page that is hosted by the author or the author's institution and is dedicated to the author's personal research interests and publication history. An author's profile page on a social media site or scholarly collaboration network site is *not* considered a personal web page. (See also: Scholarly Collaboration Network; Employer's web page).

Peer X-Press: A web-based manuscript submission system by which authors submit their manuscripts to AIP Publishing for publication, communicate with the editorial offices, and track the status of their submissions. The Peer X-Press system provides a fully electronic means of completing the License to Publish. A hard copy of the Agreement will be supplied by the editorial office if the author is unable to complete the electronic version of the form. (Conference Proceedings authors will continue to submit their manuscripts and forms directly to the Conference Editors.)

Preprint: A version of an author's manuscript intended for publication but that has not been peer reviewed and does not reflect any editorial input or publisher enhancements.

Professional Publication Ethics: AIP Publishing provides information on what it expects from authors in its "Statement of ethics and responsibilities of authors submitting to AIP Publishing journals" (<http://publishing.aip.org/authors/ethics>). AIP Publishing is also a member of the Committee on Publication Ethics (COPE) (<http://publicationethics.org/>), which provides numerous resources and guidelines for authors, editors, and publishers with regard to ethical standards and accepted practices in scientific publishing.

Scholarly Collaboration Network (SCN): Professional networking sites that facilitate collaboration among researchers as well as the sharing of data, results, and publications. SCNs include sites such as Academia.edu, ResearchGate, and Mendeley, among others.

Supplementary Material: Related material that has been judged by peer review as being relevant to the understanding of the article but that may be too lengthy or of too limited interest for inclusion in the article itself. Supplementary Material may include data tables or sets, appendixes, movie or audio clips, or other multimedia files.

U.S. Government employees: Authors working at Government organizations who author works as part of their official duties and who are not able to license rights to the Work, since no copyright exists. Government works are in the public domain within the United States.

Version of Record (VOR): The final published version of the article as it appears in the printed journal/proceedings or on the Scitation website. It incorporates all editorial input, is formatted in the publisher's standard style, and is usually viewed in PDF form.

Waiver: A request made to a university or institution to exempt an article from its open-access policy requirements. For example, a conflict will exist with any policy that requires the author to grant a nonexclusive license to the university or institution that enables it to license the Work to others. In all such cases, the Author must obtain a waiver, which shall be included in the manuscript file.

Work: The "Work" is considered all the material that comprises the article, including but not limited to the abstract, tables, figures, images, multimedia files that are directly embedded within the text, and the text itself. The Work does not include the Supplementary Material (see Supplementary Material above).

Work Made for Hire: Under copyright law, a work prepared by an employee within the scope of employment, or a work that has been specially ordered or commissioned for which the parties have agreed in writing to consider as a Work Made for Hire. The hiring party or employer is considered the author and owner of the copyright, not the person who creates the work.